

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
13 February 2003 (13.02.2003)

PCT

(10) International Publication Number
WO 03/011333 A1

(51) International Patent Classification⁷: A61K 39/10,
A01K 31/18

(21) International Application Number: PCT/US02/23670

(22) International Filing Date: 25 July 2002 (25.07.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/308,205 27 July 2001 (27.07.2001) US

(71) Applicant (for all designated States except US): BOARD
OF SUPERVISORS OF LOUISIANA STATE UNI-
VERSITY AND AGRICULTURAL AND MECHAN-
ICAL COLLEGE [US/US]; 115 System Building, 3810
W. Lakeshore Drive, Baton Rouge, LA 70803 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SUSKIND, Dana,
L. [US/US]; Apt 3, 5336 South University Avenue,
Chicago, IL 60615 (US). TILTON, Ann, H. [US/US]; 39
Pelham Avenue, Metairie, LA 70005 (US). CAIRE, Mary,
L. [US/US]; 4253 Potomac, Dallas, TX 75205 (US).

(74) Agent: RUNNELS, John, H.; Taylor, Porter, Brooks &
Phillips, L.L.P., P.O. Box 2471, Baton Rouge, LA 70821-
2471 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: BOTULINUM TOXIN IN THE TREATMENT OR PREVENTION OF ACNE

(57) Abstract: Botulinum toxin may be used to inhibit the cascade of events leading to acne. Results in preliminary studies have been dramatic. Without wishing to be bound by this theory, it is believed that botulinum toxin achieves this result through parasympathetic effects, inhibiting sweat gland activity, stimulating keratinocyte locomotion, anti-inflammatory effects, and possibly anti-androgenic effects. Botulinum toxin can play an important role in decreasing and even preventing the formation of acne.



WO 03/011333 A1

BOTULINUM TOXIN IN THE TREATMENT OR PREVENTION OF ACNE

[0001] The benefit of the 27 July 2001 filing date of United States provisional patent application serial number 60/308,205 is claimed under 35 U.S.C. § 119(e) in the United States, and is claimed under applicable treaties and conventions in all countries.

TECHNICAL FIELD

[0002] This invention pertains to a method for the treatment or prevention of acne vulgaris, particularly to the use of botulinum toxin to treat or prevent acne.

BACKGROUND ART

[0003] *Acne vulgaris*

[0004] While not life-threatening, acne vulgaris can cause significant problems for affected individuals. Depending on its severity and other factors, recalcitrant acne can be psychologically debilitating, and can impose significant financial and emotional costs on those whom it afflicts. Despite some recent successes in acne therapy, treatment failures are still common, especially in adult women. While many adults "outgrow" this disease, there are some who continue to be afflicted during much of adulthood, despite continued medical advances. Unfortunately, the most potent acne medication in current use is administered systemically via a treatment that is teratogenic, an important issue

for many women. There is an unfilled need for a more localized and effective treatment for acne, one with minimal side effects.

[0005] Acne, which most commonly occurs during adolescence, is influenced by several factors. The pathology centers on the pilosebaceous follicle, comprising the sebaceous gland, the follicle (pore), and the vellus hair. Factors that promote the formation of comedones (whiteheads or blackheads) include the following: (1) increased sebum production, (2) inflammation of the dermis and follicles by inflammatory mediators, (3) hyperkeratinization and obstruction of the upper region of the follicle, and (4) colonization of the follicle by the bacterium *Propionibacterium acnes*,

[0006] Adolescence is marked by an increase in levels of circulating androgens, particularly dehydroepiandrosterone sulfate (DHEAS). The increased androgen levels are thought to cause sebaceous glands to enlarge and to increase sebum production. While most acne patients have normal hormone levels, there are reasons to conclude that increased sebum production plays an important role in acne. For example, there is a correlation between the rate of sebum production and the severity of acne. In addition, acne patients typically produce sebum that is deficient in linoleic acid, which is a potential cause of abnormal keratinization and follicular obstruction.

[0007] In response to increased sebum levels, *Propionibacterium acnes*, a gram positive, anaerobic, diphtheroid bacterium, often colonizes the sebaceous follicles. *P. acnes* exacerbates acne by acting as a chemo-attractant for neutrophils (a type of white blood cells, also called polymorphonuclear leukocytes, or PMNs). The neutrophils ingest the *P. acnes*, and in doing so release various hydrolytic enzymes that damage the follicular wall. The released follicular contents then invade the dermis and cause an inflammatory reaction, manifesting as pustules, erythematous papules, or nodules. In a separate route, *P. acnes* can hydrolyze triglycerides to free fatty acids, which also increase inflammation and follicular obstruction. *P. acnes* may also activate the complement components of the immune system, which can also lead to follicular obstruction.

[0008] The follicles are lined with squamous epithelium, a layer of cells that is contiguous with the skin surface. In an acne-prone individual, the

shedding of cells from this lining is often impeded, perhaps due to an increased level of intracellular adhesion that promotes the retention of cells. The retained cells can obstruct the follicles, resulting in comedones. The exact cause of this inhibited shedding is unknown, but it may be related either to abnormalities in epidermal differentiation, or to abnormal sebum composition (e.g., a deficiency in linoleic acid).

[0009] It has also been demonstrated that increased sebum levels can irritate keratinocytes, causing the release of interleukin-1, which in turn can cause follicular hyperkeratinization.

[0010] The final common pathway in each of these acne-causing routes, which are not mutually exclusive, is follicular obstruction.

[0011] ***Current Acne Therapies***

[0012] Currently-used acne therapies are directed at various aspects of the acne cascade. The most commonly used therapy is probably topical benzoyl peroxide, which has an antibacterial effect, and which may also decrease free fatty acids, resulting in a decrease in inflammation and follicular obstruction.

[0013] Topical and systemic antibiotics have been used to target *P. acnes*. The antibiotics that have been used for this purpose include erythromycin, tetracycline, clindamycin, and doxycycline. Predictably, the prolonged use of antibiotics often leads to the development of resistant strains of *P. acnes*.

[0014] Topical and systemic retinoids (derivatives of vitamin A) have been used to normalize the keratinization of the follicle, decreasing follicular obstruction and rupture. Systemic isotretinoin (Accutane™), which is highly effective for acne, unfortunately has serious adverse effects, most notably teratogenicity. Topical retinoids include tretinoin (Retin-A™), whose chemical structure is similar to that of isotretinoin.

[0015] The current state of the art in treating acne is summarized, for example, in the following two review articles: D. Krowchuk, "Managing Acne in Adolescents," *Pediatric Dermatology*, vol. 47, pp. 841-857 (2000); and B. Johnson *et al.*, "Use of Systemic Agents in the Treatment of Acne Vulgaris," *American Family Physician*, vol. 62, pp. 1823-1830, 1835-1836 (2000).

[0016] Advances in acne therapy have followed better understanding of its multiple causes. Most treatments are directed at normalizing keratin production (e.g., through the use of retinoids), or at controlling bacterial colonization. To the inventors' knowledge, no prior treatments have sought to affect either sweat gland production or keratinocyte locomotion, both of which can be factors in the follicular occlusion that leads to acne.

[0017] *Botulinum toxins and Pharmacology*

[0018] *Clostridium botulinum*, an anaerobic bacterium, produces seven toxins that have similar neurotoxic effects, but that are antigenically distinct: serotypes A through G. These toxins are potent neuromuscular agents that inhibit the release of acetylcholine at neuromuscular junctions and at neuroglandular junctions. See M.F. Brin, "Botulinum toxin: Chemistry, Pharmacology, Toxicity, and Immunology," *Muscle and Nerve Supp.* 6, pp. S146-S168 (1997); and I. Kinkelin *et al.*, "Effective Treatment of Frontal Hyperhidrosis with botulinum toxin A," *Brit. J. Dermatol.*, vol. 143, pp. 824-827 (2000).

[0019] Neither the synthesis nor the storage of acetylcholine is affected by botulinum toxin. Therefore, its effects are temporary. Botulinum toxin penetrates the endosomal membrane into the cytosol, where the secretion of acetylcholine is blocked. Botulinum toxin also appears to possess anti-inflammatory effects. Studies of the effect of botulinum toxin on glandular function have observed that its effects occur at the presynaptic terminal of the parasympathetic cholinergic nerves.

[0020] Botulinum toxin A (sold under the trademark Botox® by Allergan (Irvine, California), and under the trademark Dysport® by Ipsen Limited (Maidenhead, Berkshire, United Kingdom)) has been used for certain therapeutic purposes at both neuroglandular and neuromuscular junctions. Its temporary neuromuscular blockade has found numerous previous uses, including the treatment of strabismus, migraines, achalasia, hemifacial spasm, rhytids, and spastic dysphonia. The injections have proven to be safe and effective for these purposes, with only minor side effects such as local swelling and transient weakness of nearby muscles. Botulinum toxins A and B have been reported to be well tolerated in patients who have used it.

[0021] Intracutaneous botulinum toxin A has become a primary therapy for axillary, palmar, solar, and frontal hyperhidrosis (excessive sweating), as well as for gustatory sweating (Frey's Syndrome). It has also been shown to be an effective alternative to surgery in Hailey-Hailey disease, a benign familial pemphigus that affects primarily the groin and axillary folds (areas of high moisture).

[0022] Studies have found no long-term effects of botulinum toxin A in human tissue. Histological studies have found no nerve fiber degeneration, and no sweat gland atrophy as a result of therapy. Antibodies to botulinum toxin A have been found in fewer than 5% of patients receiving botulinum toxin A injections. There have been no reported cases of anaphylaxis in response to injections of botulinum toxin A.

[0023] Botulinum toxin B (sold under the trademark Myobloc™ and NeuroBloc™ by Elan Pharmaceuticals (Dublin, Ireland)) has been approved by FDA for use in treating patients with cervical dystonia.

[0024] All seven botulinum toxin serotypes (A through G) are available commercially from Metabionics, Inc. (Madison, Wisconsin).

[0025] *Neuromuscular Junction Blockade*

[0026] The first reported clinical use of botulinum toxin A was its use in the treatment of strabismus in the 1970's. Injecting the small extraocular muscles resulted in a realignment of the muscles, straightening of the globe, and improvement of visual alignment in some cases. Subsequently, botulinum toxin A has been used therapeutically for several other purposes, both functional and cosmetic. It has proven effective in treating focal dystonias, spasmodic dysphonia, blepharospasm, hemifacial spasm, torticollis, and cervical dystonia. It has also recently been used in facial plastic surgery, especially to treat rhytids (wrinkles).

[0027] In 1990, the use of botulinum toxin A was reported in ambulatory and nonambulatory cerebral palsy patients, namely by injecting different muscle groups to treat spastic diplegia. For example, double-blinded, placebo-controlled studies on the effect of botulinum toxin A on limb dystonias and spasticity have found significant subjective as well as objective benefits.

[0028] The American Academy of Ophthalmology, the American Academy of Neurology, the American Academy of Otolaryngology, and the National Institutes of Health have all released statements supporting the therapeutic efficacy of botulinum toxin A for a variety of clinical conditions. Among the conditions for which botulinum toxin A has been used are blepharospasm, strabismus, cervical dystonia, spasmodic torticollis, rhytids, hemifacial spasm, facial spasm, spasmodic dysphonia, focal hand dystonia, hyperfunctional facial wrinkles, Frey's syndrome, hyperhidrosis, adult spasticity, adjunctive treatment of spasticity in cerebral palsy, oromandibular dystonia, and other dystonias.

[0029] *Neuroglandular Junction Blockade*

[0030] The effects of botulinum toxin A as an anticholinergic agent at the neuroglandular junction have not been explored as extensively as those occurring at the neuromuscular junction. Clinical studies examining the effect of

intracutaneous botulinum toxin for focal hyperhidrosis found complete abolition of sweating in the injected area within 3 to 7 days. No adverse effects were reported, and in a five month follow-up there were no clinical recurrences of the hyperhidrosis. See generally I. Kinkelin *et al.*, "Effective Treatment of Frontal Hyperhidrosis with botulinum toxin A," *Brit. J. Dermatol.*, vol. 143, pp. 824-827 (2000).

[0031] Gustatory sweating is another area of neuroglandular dysfunction in which botulinum toxin A has proven effective. Gustatory sweating (or Frey's syndrome) is a disabling disorder in which the cheek skin sweats profusely during eating. The syndrome often occurs after parotidectomy, and may be due to the misdirection of the regenerating parasympathetic fibers that enervate the sweat glands of the face. Intracutaneous botulinum toxin A has been reported to significantly decrease or prevent sweating for over six months, with no clinical evidence of facial weakness in any patients.

[0032] Botulinum toxin A injected into the submandibular glands has been reported to significantly decrease salivation resulting from stimulation of the lingual nerves. The decreased salivation was temporary, and did not appear to be directly toxic to the acinar cells of the gland. See D. Suskind *et al.*, "Clinical study of botulinum A toxin in the treatment of sialorrhea in children with cerebral palsy," *Laryngoscope*, vol. 112, pp. 73-81 (2002).

[0033] Canine studies have also shown that vasomotor rhinorrhea, a parasympathetically controlled phenomenon, responds to topical botulinum toxin A.

[0034] While the duration of botulinum toxin A's action at the neuromuscular junction appears to be about three months, there appears to be a longer-lasting effect at the glandular level. Botulinum toxin A has produced anhydrosis for over 12 months in patients with gustatory sweating. The reason for the difference in duration of action is uncertain. Hypotheses include a higher rate of re-synthesis of SNAP-25 (the protein cleaved by botulinum toxin) in

neuromuscular synapses, and a higher area of axonal sprouting and consecutive reinnervation of muscle fibers as compared to that in glandular tissue.

[0035] To the inventors' knowledge, there have been no prior suggestions that botulinum toxin might be used in the treatment or prevention of acne.

DISCLOSURE OF THE INVENTION

[0036] We have discovered that botulinum toxin may be used to inhibit the cascade of events leading to acne. Results in preliminary studies with botulinum toxin A have been dramatic. Without wishing to be bound by this theory, it is believed that botulinum toxin achieves this result through parasympathetic effects, inhibiting sweat gland activity, and stimulating keratinocyte locomotion. Associated anti-inflammatory and anti-androgenic effects may also contribute. Treatments may be repeated periodically as needed to inhibit the recurrence of acne, typically at intervals between about 3 months and about 6 months, preferably about once every 4 months.

[0037] Without wishing to be bound by this theory, we believe that botulinum toxin inhibits the formation of acne through at least three different pathways. First, botulinum toxin inhibits sweat production by sweat glands. Decreased perspiration may clinically improve acne by decreasing the growth of *P. acnes*. Thus the ability of botulinum toxin to decrease perspiration leads to decreased acne. Second, follicular occlusion by keratinocytes is the final common pathway in each of the various routes leading to acne. Keratinocyte locomotion is inhibited by the high dose stimulation of nicotinic acetylcholine receptors. By inhibiting the release of acetylcholine, botulinum toxin may indirectly increase the migration of keratinocytes, thus reducing follicular occlusion. Third, the androgen surge during puberty is a known instigator of acne, and studies have shown that androgens increase the number of acetylcholine receptors. Interestingly, androgen receptors are found on pilosebaceous duct keratinocytes, which are important in follicular occlusion. Thus we believe that during puberty androgens increase the number of

acetylcholine receptors on the pilosebaceous keratinocytes, leading to further inhibition of keratinocyte locomotion through increased acetylcholine stimulation. By inhibiting the release of acetylcholine, botulinum toxin decreases the number of acetylcholine receptors on the pilosebaceous keratinocytes, thereby increasing keratinocyte locomotion through decreased acetylcholine stimulation. Additionally, botulinum toxin may have anti-inflammatory effects.

MODES FOR CARRYING OUT THE INVENTION

[0038] Example 1

[0039] The "index case" was a Caucasian woman in her late twenties who had been afflicted by severe cystic acne since puberty. She experienced a complete resolution of her acne following intracutaneous botulinum toxin A injections. An otherwise healthy woman, she had tried several prior medical treatments for her acne, including both spironolactone and three courses of isotretinoin, without resolution.

[0040] Following initial injections of botulinum toxin A, her forehead, which had previously been covered with deep cystic lesions, completely resolved.

[0041] She now receives botulinum toxin A injections in her forehead, chin, and cheek/nasolabial fold area approximately once every six months. After about six months, the cysts begin to recur. They resolve about one week after injections. Injections are spaced approximately 1.5 cm apart from one another, and the dose per injection site is about 2.5 U.

[0042] Her physical examination now demonstrates an attractive blonde woman with strikingly clear skin. Her facial nerve is intact, with only slight to moderate weakness of her frontalis muscles. She states that the botulinum toxin A injections and the resulting resolution of her acne have had a profound effect upon her life, both socially and psychologically.

[0043] A dermatologist has confirmed the substantial improvement in this patient's condition.

[0044] Example 2

[0045] A Caucasian woman in her early thirties presented with severe cystic acne, which she had since puberty, and which had not responded to traditional treatments. She already had severe scarring as a result of her acne. Injections of botulinum toxin A were made in her back. Within one week there was a substantial decrease in acne in the injected region, and by one month there a significant decrease in the papules and pustules. A discernable demarcation line was visible where the injections ended, as there were active lesions around the periphery, beginning about 1 cm from the zone where the injections had been made. A total of 86 units of Botulinum toxin A were used, 2.5 units per injection at injection sites spaced about 2-3 cm from one another.

[0046] A dermatologist has confirmed the substantial improvement in this patient's condition. The improvement lasted for about five months prior to the return of pruritus and papules.

[0047] Example 3

[0048] A 20-year-old Caucasian man who had severe cystic acne since puberty presented with acne involving his face, neck, chest, and back. He had previously undergone two courses of isotretinoin, as well as many courses of systemic and topical antibiotic treatments, all without resolution of his acne. His only other significant medical problem was Crohn's colitis. A total of 162 U of botulinum toxin A was injected into his back, but not the other areas where acne was present. There was a substantial clearing of the treated areas after one week, and almost complete clearing after three weeks. By contrast, there was no improvement in the untreated areas (i.e., face, neck, and chest).

[0049] A dermatologist has confirmed the substantial improvement in this patient's condition. The clearing persisted for five months during follow-up observations.

[0050] Although the number of patients in whom the botulinum toxin treatment has been tested in this initial experiments was small -- only three patients -- dramatic improvements were seen in 100% of these initial patients, all of whom had recalcitrant cases of acne that had not responded to conventional

treatments. A subsequent, small-scale clinical study has now been conducted as well, with the results reported below.

[0051]

Example 4

[0052] A small-scale clinical study is being conducted to examine the clinical benefit of intradermal botulinum toxin A (Botox™) injections in the prevention or reduction of acne vulgaris lesions in adults. As of the filing date of this application, this study is ongoing, but some preliminary results were available. The protocol being used in this study is as follows:

[0053] The study is an open-label, prospective, single-center, pilot study. Fifteen subjects are to begin participation in the study. Informed consent is obtained from all subjects.

[0054] ***Eligibility criteria***

Eligibility criteria are as follows:

[0055] **Inclusion**

1. Males or females.
2. General good health as determined by the initial paperwork.
3. Willingness to have clinical examinations and photographs of the face and back.
4. Diagnosis of recalcitrant acne (unresponsive to conventional therapy, including systemic antibiotics).
5. Negative pregnancy test (females).

6. At least 10 of the 15 subjects enrolled into the study will have cystic acne on the forehead or back (at least half must be forehead).
7. Up to 5 subjects will have non-cystic inflammatory acne on the forehead or back (at least 3 must be forehead).
8. Individuals must be willing to serve as a no-treatment control group. Three of the 15 subjects were assigned to the no treatment group (at least 2 from the cystic group and 1 from the non-cystic group).
9. Able to understand the requirements of the study and sign an informed consent form.
10. Willingness to remove facial makeup no less than 30 minutes prior to each facial exam.

[0056] Exclusion

1. Females who had begun treatment with estrogens or birth control pills for 12 weeks or less prior to baseline.
2. Subjects with any skin disease that might interfere with the diagnosis or evaluation of acne.
3. Subjects currently participating in an investigational drug study or who had participated in one within 30 days of the baseline visit.
4. Individuals with uncontrolled metabolic disease such as diabetes, hypertension, hyperthyroidism, or hypothyroidism as determined by the health questionnaire.
5. Women known to be pregnant, nursing, or planning to become pregnant within the next six (6) months as determined by the initial paperwork.

6. Individuals having an allergy or sensitivity to botulinum toxin A.
7. Individuals with a history of poor cooperation, non-compliance with treatment, or unreliability in previous clinical studies.
8. Individuals with a history of Myasthenia Gravis or Eaton Lambert Disease.
9. Individuals who had a change in antibiotic use within the last 1.5 months.
10. Individuals who were receiving aminoglycosides, anticholinesterases, magnesium sulfate and other drugs or products that interfere with neuromuscular transmission.
11. Individuals who have had treatments with or exposure to botulinum toxin A.

[0057] Individuals are admitted to study at the discretion of the investigator based on medical history and findings of the pre-study interview and examination. Each subject is expected to complete the full course of the study.

[0058] CONDUCT OF STUDY

Procedures

[0059] Visit 1 – Screening

Prospective subjects complete a health/eligibility questionnaire, a photography release form, a confidentiality agreement, and an informed consent agreement. Those who qualify are examined on the face and back for the presence of cystic or inflammatory acne.

[0060] Fifteen qualified subjects, as described in the eligibility criteria, enroll in the study and return for baseline in 1-3 days.

[0061] Subjects are placed into a treatment group based on the location of their acne (forehead or back). Each subject is treated either on the forehead or

back, but not both. Randomization of treatment groups occurs between Visits 1 and 2 (after all individuals are screened and all subjects are enrolled).

[0062] Visit 2 - Baseline

Subjects return to the clinic 1-3 days after the qualification visit. Subjects who in both the treatment group and the control group completed a pre-injection questionnaire.

[0063] A trained clinical grader counts papules, pustules, open comedones, closed comedones, and cysts at the test site. The grader also assesses erythema and elevation of the lesions, as well as dryness and oiliness of the surrounding skin. A single cross-polarized photograph is taken of the test site.

[0064] A topical anesthetic cream is applied to the test sites prior to injections. The study investigator performs Botox™ injections in the forehead with a distance between adjacent injection points ranging from 1.5 to 2 cm. Control patients were injected with saline only. No injections are performed closer than 1.5 cm to the brow to prevent ptosis. The intracutaneous nature of the injections is expected to decrease the likelihood of muscle weakness (as compared to botulinum toxin injections for treating wrinkles. Injections on the back are conducted on a similar spacing, depending on the individual's acne distribution. Approximately 2.5 - 3 mouse units of Botox™ are injected into each area, as otherwise described in M. Naumann *et al.*, "Focal Hyperhidrosis: Effective Treatment with Intracutaneous Botulism Toxin," *Arch. Dermatol.*, vol.134, pp. 301-304 (1998); and M. Naumann *et al.*, "Treatment of Gustatory Sweating with Botulism Toxin," *Ann. Neurol.*, vol. 42, pp. 973-975 (1997). The dilution is 3 units per 0.05 cc, equivalent to 1.66 cc per 100 units of botulinum toxin A. The upper dose limits are 80 units for the forehead, and 125 units for the back. Subjects remain in the clinic for 30 minutes after receiving injections. The subjects' current acne regimens are continued.

[0065] Subjects are given a calendar of future visits. A clinician reviews study instructions with each subject, and gives them a written copy for use at home.

[0066] Visits 3-7 - Further Evaluations

Subjects return to the clinic approximately 1, 2, 4, 8 and 12 weeks after treatment (Visits 3, 4, 5, 6 and 7, respectively). Evaluations are performed at each visit as described for baseline, but no further injections are administered. On each visit, subjects complete a questionnaire; and a trained clinical grader performs acne lesion counts, grades lesions for erythema and elevation, and grades surrounding skin for dryness and oiliness. A single cross-polarized photograph is repeated at each test site.

[0067] Any adverse events are noted.

[0068] If a subject fails to return for a scheduled examination, a staff member attempts to contact the subject to determine whether the subject has continued to follow study instructions and intends to continue participation in the study. The subject is processed as soon as possible, or is allowed to miss a study visit at the discretion of the investigator. Subjects lost from the study are documented.

[0069] Examples 5-9**[0070] Preliminary Results**

[0071] As previously mentioned, this study is ongoing as of the filing date of this application. Preliminary results from a portion of the study that has been completed are reported below.

[0072] Results after 12 weeks for the 5 subjects in the "back" group are summarized in Table 1.

Table 1 – “Back” Group (n = 5)

Variable	Mean Baseline Score	Mean Score after 12 weeks	Mean Change, Baseline to 12 weeks	Percent Change in Mean, Baseline to 12 weeks	Standard Error of the Mean, Baseline to 12 weeks	P (change, baseline to 12 weeks) ** = statistically significant change (P<0.05)
Papules	19.0	4.8	-14.2	-75%	5.3	0.01**
Pustules	1.8	0	-1.8	-100%	2.5	0.18
Open Comedones	5.4	1.8	-3.6	-90%	4.4	0.14
Closed Comedones	5.0	4.8	-0.2	-3%	1.8	0.81
Cysts	1.2	0.6	-0.6	-70%	0.9	0.21
Lesion Erythema	1.8	0.9	-0.9	-55%	0.6	0.02**
Lesion Elevation	1.6	0.8	-0.8	-50%	0.6	0.03**
Dryness	0.2	0.0	-0.2	-100%	0.5	0.37
Oiliness	0.2	0.2	0.0	0%	0.0	---

[0073] Although the sample size for these preliminary results was small (n = 5), statistically significant decreases were seen in the number of papules, the degree of lesion erythema, and the degree of lesion elevation from baseline to 12 weeks. Large numerical differences were seen in three other measurements, differences that were not statistically significant for this small sample size, but that may well become statistically significant as additional data become available: number of pustules, number of open comedones, and number of cysts. With the same caveat about sample size, these preliminary results suggest minimal or no differences resulting from this particular treatment in the number of closed comedones, incidence of dryness, and incidence of oiliness. No adverse events, toxicity, or side effects were reported.

[0074] Example 10 -- Dosage and Toxicity

[0075] An important consideration is the choice of proper dosage. Proper dosages will be established through otherwise standard dose-response studies and toxicity studies. Because the target area for this treatment is similar to that for hyperhidrosis, similar dosages have been chosen for initial studies. See, e.g., M. Naumann *et al.*, "Focal Hyperhidrosis: Effective Treatment with Intracutaneous Botulism Toxin," *Arch. Dermatol.*, vol. 134, pp. 301-4 (1998); and M. Naumann *et al.*, "Treatment of Gustatory Sweating with Botulism Toxin," *Ann. Neurol.* vol. 42, pp. 973-975 (1997). There appear to have been no studies defining an accurate dose-response curve for hyperhidrosis, although general clinical guidelines have been generated. See Table 2. Clinical preparations of botulinum toxin A, which is made under strict FDA supervision in the United States, are measured in mouse units. A mouse unit, 1 U, is defined as the amount of botulinum toxin A that kills fifty percent (LD_{50}) of a group of 18-20 gram female Swiss-Weber mice by parenteral injection. Contraindications to the use of botulinum toxin A are very few. (Note: There have been reported suggestions that the Botox® unit is 3-5 times as potent as the Dysport® unit. The botulinum toxin A used in the experiments reported here was obtained as Botox®; so, to the extent that there may be a difference in the potency of these two commercially available preparations of botulinum toxin A, the "units" used in the present specification and claims may be interpreted with reference to Botox® units.)

[0076]

Table 2
Human Botulinum-A Toxin
Predicted Dose-Response

Dosage Unit (U)

1 U = LD_{50} mice \approx 0.04 ng

Clinical Efficacy

1-12 U per kg body weight, depending on patient's response, routine maximum 400 U, delivered not more often than once per 90 days; lower or higher doses may be appropriate for some patients

Toxicity

Oral: 3,000-250,000 U (monkey)

Parenteral: 39 U / kg (monkey)

Assumption:

Monkey LD₅₀ = Human LD₅₀

Adult LD₅₀ = 3000 U

Safety Factor

toxicity = 40 U/kg

[0077] Based on the preliminary studies reported above, it is expected that in most cases intracutaneous injections should be spaced about 0.5 or 1 cm to about 10 cm apart from one another, preferably about 1.5 to about 3 cm apart; and that the dose per injection site should be about 1 U to about 10 U or 20 U botulinum toxin A, preferably about 2 to about 3 U. These dosages may be varied based on individual responses to the injections.

[0078] The botulinum toxin may be injected intracutaneously or subcutaneously. Alternatively, botulinum toxin may be injected intramuscularly at similar doses, particularly when it is desired to simultaneously weaken certain facial muscles, e.g., to reduce rhytids.

[0079] Alternatively, the botulinum toxin may be applied to the skin as a topical or percutaneous cream, gel, ointment, lotion, aerosol or spray, powder, paste, fixed dressing or other topical or percutaneous administration to provide similar clinical efficacy. Superficial application of the botulinum toxin may particularly help to improve comedonal acne, the inflammatory aspects of acne,

and topical application may help increase absorption. Such topical administrations are best reconstituted as needed and then stored under refrigeration, as they have a limited shelf life. The shelf life can be extended to a few weeks or longer by adjusting the pH to be slightly acidic, e.g., about pH 5.6. Topical vehicles may include other components used in the art in topical medications, such as vanishing vehicles, moisturizers, emollients, chemical carriers (e.g., DMSO), and buffers known in the art. Typical concentrations of active ingredient may be on the order of 5 to 200 U per mL, depending on the absorption rate and the toxin serotype. In general, the concentration should be adjusted so that the effective rate of administration is equivalent to that of effective injections of botulinum toxin. The use of ultrasound, or of an electrophysiological current device, such as iontophoresis, may enhance absorption of topically applied toxin. It is known in the art (albeit using other compounds for treating other conditions) that topically applied agents can reach and clinically affect the sweat glands and sebaceous glands. See, e.g., F. Marzulli *et al.* (Eds.), "Dermatotoxicology and Pharmacology" in *Advances in Modern Toxicology*, vol 4, John Wiley and Sons, New York (1977).

[0080] In preparations that are currently available commercially, botulinum toxin A is stored frozen; while botulinum toxin B is stored at room temperature.

[0081] Examples 11 and 12

[0082] Preliminary trials have been conducted on topical applications of botulinum A toxin with two acne patients. Additional, larger-scale trials of topical applications will be conducted to confirm what types of acne respond well to topical treatment. For the topical treatment, botulinum toxin A was reconstituted at a concentration of 100 Units per 0.5 mL of solvent. For the mid-back treatments, the solvent was a 1.4% aqueous solution of polyvinyl alcohol as a neutral ophthalmic solution, and for the left back treatments the solvent was preservative-free saline. A 0.2 mL or 40 U portion of the solution was applied to

an area 6 cm x 4 cm, patched with Tegaderm™ for 4 hours to avoid disturbances. Results are presented below.

[0083]

Table 3

	Patient 1 Baseline	Patient 1 Week 2	Patient 1 Week 4	Patient 2 Baseline	Patient 2 Week 2	Patient 2 Week 4
Lesions, Mid-Back						
Papules	5	1	0	1	10	2
Pustules	0	0	0	0	0	0
Open Comedones	4	1	1	0	0	0
Closed Comedones	0	2	1	11	0	4
Nodules	0	0	0	0	0	0
Lesions, Left Back						
Papules	5	3	3	10	5	3
Pustules	2	0	0	0	0	0
Open Comedones	0	0	0	0	0	0
Closed Comedones	0	6	2	4	1	2
Nodules	0	0	0	0	0	0

[0084] There is a potential emerging trend in these data towards the improvement of acne following the topical application of botulinum toxin. Further studies are ongoing. Due to the small number of patients in these initial experiments on topical application, a statistical analysis at this time would have little meaning. Topical application of the botulinum toxin to treat acne may eventually prove to be the preferred method of application, both because of the

ease of application, and because of the ease of directly delivering the medication directly to affected areas where there is inflammation or other disruption of the skin's tight junctions.

[0085] Example 13

[0086] Further Phase II Clinical Trials will be conducted to confirm that intracutaneous or topical botulinum toxin A injections can safely and effectively prevent or decrease the incidence of recalcitrant acne in adults and adolescents. Following Phase II trials, Phase III trials will be conducted, all in accordance with applicable laws and regulations.

[0087] The inventors speculate that it is possible that a subpopulation of patients who have previously been treated with botulinum toxin for other reasons (e.g., rhytids) may have also incidentally experienced improvements in acne as a side effect. However, to the inventors' knowledge, no previous report has ever suggested the presence of such a side effect of botulinum toxin treatment, nor has any previous report suggested that there should be any causal relation between administration of botulinum toxin and improvement in acne.

[0088] As used in the specification and claims, an "effective amount" of botulinum toxin is an amount sufficient to prevent acne, or to inhibit the formation of acne, or to reduce the level of acne, to a clinically significant degree. "Significance" for this purpose is determined as the $P < 0.05$ level, or by such other measure of statistical significance as is commonly used in the art for a particular type of experimental determination.

[0089] While the preliminary experiments reported above used botulinum toxin A, without wishing to be bound by this theory, it is believed that the other botulinum toxin serotypes, namely botulinum toxins B, C, D, E, F, and G, may also be used in the method of this invention to treat or prevent acne. Proper doses and toxicity will be determined as generally described above for botulinum toxin A. There have been suggestions that botulinum toxin C may have two variants, one with neurotoxic activity and one lacking such activity. To the extent

that such variants of any of the botulinum toxin serotypes do exist, the use of all such variants displaying anti-acne activity is considered to be within the scope of this invention.

[0090] For example, changing the serotype used might be indicated if a patient develops an immune response to a particular serotype. The dosages and durations of effect may differ among the different serotypes, as will be determined by experiments generally along the lines of those discussed above for serotype A. To reduce the likelihood of developing an immune response, the smallest effective dose for each patient should be used, and the time between successive treatments should be made as long as possible without having substantial symptoms return.

[0091] The complete disclosures of all references cited in this specification are hereby incorporated by reference. In the event of an otherwise irreconcilable conflict, however, the present specification shall control.

What is claimed:

1. A method for inhibiting acne vulgaris in a human who is susceptible to acne vulgaris, said method comprising delivering an effective amount of botulinum toxin A to one or more regions of the skin of the human that are susceptible to acne vulgaris.
2. A method as recited in Claim 1, wherein said delivering comprises the intracutaneous injection of botulinum toxin A at multiple sites in the skin, wherein the sites of adjacent injections are separated by about 0.5 to 10 cm.
3. A method as recited in Claim 1, wherein said delivering comprises the intracutaneous injection of botulinum toxin A at multiple sites in the skin, wherein the sites of adjacent injections are separated by about 1.5 to 3 cm.
4. A method as recited in Claim 1, wherein said delivering comprises the topical application to susceptible regions of the skin of a composition comprising botulinum toxin A.
5. A method as recited in Claim 1, wherein said delivering comprises the subcutaneous injection of the botulinum toxin A.
6. A method as recited in Claim 1, wherein said delivering comprises the intra-muscular injection of the botulinum toxin A.
7. A method as recited in Claim 1, wherein said method is repeated periodically to inhibit the recurrence of acne vulgaris.
8. A method as recited in Claim 1, wherein said method is repeated at intervals from about 3 months to about 6 months to inhibit the recurrence of acne vulgaris.

9. A method as recited in Claim 1, wherein said method is repeated at intervals of about 4 months to inhibit the recurrence of acne vulgaris.

10. A method as recited in Claim 1, wherein said delivering comprises the intracutaneous injection of botulinum toxin A at multiple sites in the skin, and wherein between about 1 U and about 20 U of botulinum toxin A is injected at each site.

11. A method as recited in Claim 1, wherein said delivering comprises the intracutaneous injection of botulinum toxin A at multiple sites in the skin, and wherein between about 2 U and about 3 U of botulinum toxin A is injected at each site.

12. A method as recited in Claim 1, additionally comprising the inhibition of at least one rhytid by said delivering of an effective amount of botulinum toxin A to one or more regions of the skin that are susceptible to acne vulgaris and that also contain at least one rhytid.

13. A method for inhibiting acne vulgaris in a human who is susceptible to acne vulgaris, said method comprising delivering an effective amount of botulinum toxin to one or more regions of the skin of the human that are susceptible to acne vulgaris.

14. A method as recited in Claim 13, wherein said delivering comprises the intracutaneous injection of botulinum toxin at multiple sites in the skin, wherein the sites of adjacent injections are separated by about 0.5 to 10 cm.

15. A method as recited in Claim 13, wherein said delivering comprises the intracutaneous injection of botulinum toxin at multiple sites in the skin, wherein the sites of adjacent injections are separated by about 1.5 to 3 cm.

16. A method as recited in Claim 13, wherein said delivering comprises the topical application to susceptible regions of the skin of a composition comprising botulinum toxin.

17. A method as recited in Claim 13, wherein said delivering comprises the subcutaneous injection of the botulinum toxin.

18. A method as recited in Claim 13, wherein said delivering comprises the intra-muscular injection of the botulinum toxin.

19. A method as recited in Claim 13, wherein said method is repeated periodically to inhibit the recurrence of acne vulgaris.

20. A method as recited in Claim 13, wherein said method is repeated at intervals from about 3 months to about 6 months to inhibit the recurrence of acne vulgaris.

21. A method as recited in Claim 13, wherein said method is repeated at intervals of about 4 months to inhibit the recurrence of acne vulgaris.

22. A method as recited in Claim 13, additionally comprising the inhibition of at least one rhytid by said delivering of an effective amount of botulinum toxin to one or more regions of the skin that are susceptible to acne vulgaris and that also contain at least one rhytid.

23. A method as recited in Claim 13, wherein the botulinum toxin comprises botulinum toxin B.

24. A method as recited in Claim 13, wherein the botulinum toxin comprises botulinum toxin C.

25. A method as recited in Claim 13, wherein the botulinum toxin comprises botulinum toxin D.

26. A method as recited in Claim 13, wherein the botulinum toxin comprises botulinum toxin E.

27. A method as recited in Claim 13, wherein the botulinum toxin comprises botulinum toxin F.

28. A method as recited in Claim 13, wherein the botulinum toxin comprises botulinum toxin G.

29. A method as recited in Claim 13, additionally comprising the inhibition of at least one rhytid by said delivering of an effective amount of a botulinum toxin to one or more regions of the skin that are susceptible to acne vulgaris and that also contain at least one rhytid; wherein the botulinum toxin is selected from the group consisting of botulinum toxin B, botulinum toxin C, botulinum toxin D, botulinum toxin E, botulinum toxin F, and botulinum toxin G.

INTERNATIONAL SEARCH REPORT

 International application No.
 PCT/US02/29670

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : A61K 39/10; A01K 31/16 US CL : 424/239.1; 514/2 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/239.1; 514/2 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet.		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,670,484 A1 (BINDER) 23 September 1997, see abstract, col. 1, lines 13-14, and 17; col. 1, lines 35-61; col. 2, lines 19-23; col. 2, line 45; col. 3, lines 23-46; col. 4, lines 62-65; col. 5, lines 7-26, col. 6, lines 10-37; Table 1, claims 1-3, and entire document.	1-7,10-19, 22-29 ----- 8-9 20-21
Y		
Y,P	US 6,447,787 A1 (GASSNER et al), 10 September 2002, see all claims, col. 1, lines 43-56 and col. 2, lines 1-16, and entire document.	1-29
A	US 5,766,605 A1 (SANDERS et al), 16 June 1998, see claims abstract and entire document.	1-29
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents:		
"A"	document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier document published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	
"P"	document published prior to the international filing date but later than the priority date claimed	"A" document member of the same patent family
Date of the actual completion of the international search 05 OCTOBER 2002		Date of mailing of the international search report 23 DEC 2002
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer Valérie Bell-Harris for GINNY FORTNER Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/23670

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ROENIGK, H.H., JR., Combination resurfacing, Dermatologic therapy, 2000, Vol. 13, No. 2, pages 215-222, (abstract only)	1, 4

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US02/23670

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

WEST, DIALOG

search terms: acne, vulgaris, botulinum, botox, botx, btna, btxa , dermatitis, skin, epidermis, pustula, follicle